

First report of Amblypygi (Charinidae: *Charinus ioanniticus*) from Turkey

První zpráva o bičovci *Charinus ioanniticus* (Amblypygi: Charinidae) z Turecka

František KOVÁŘÍK¹⁾ & Daniel VLASTA²⁾

¹⁾U Botiče 1/1389, CZ-140 00 Praha 4, Czech Republic

²⁾Vysočanská 546, CZ-190 00 Praha 9, Czech Republic

Faunistics, taxonomy, Amblypygi, Charinidae, *Charinus ioanniticus*, Turkey

Abstract. *Charinus ioanniticus* (Kritscher, 1959) is recorded for the first time from Turkey. All ten specimens collected were found at night in a karst cave at the outskirts of the Çevlik village near Samandagi.

Charinus ioanniticus (Kritscher, 1959)

Lindosiella ioannitica Kritscher, 1959: 454; Kraus, 1961: 491.

Charinus sp. ?: Rosin & Shulov, 1960: 167-168.

Charinus ioanniticus: Weygoldt, 1972: 123.

Material. Turkey, Çevlik near Samandagi, 2 females, 23.v.1995, Vladislav Šejna leg., coll. Department of Invertebrate Zoology, National Museum (Natural History), Prague; 1 male, 1 female, 6 juvs, 27.-28.ix.1995, Dan Vlasta leg., in the author's collections.

Comments. All specimens were found at night in a single 30 m long karst cave at the outskirts of the Çevlik village. The vicinity of the cave is intensely utilised for agricultural purposes. The cave entrance faces the sea which is about 400 m away. The more spacious front part of the cave occasionally serves as a shelter for herdsmen and their animals, whereas the narrow back part has undisturbed speleothems and is inhabited by bats. The floor is covered by a thin layer of silty earth with embedded limestone boulders of varying size and fragments of fallen speleothems. Water seeps into the cave through wall fractures and saturates much of the floor deposit. The fractures probably connect the cave with a large, inaccessible karst system.

Distribution. *Charinus ioanniticus* was first found on the Greek island of Rhodos (Kritscher 1959: 457) and a year later also in Israel, Jerusalem (Rosin & Shulov 1960: 167). However, Kraus (1961: 491) found one female in Zipari near Nazareth more than a decade earlier, on 5.viii.1948. Weygoldt (1972: 129) listed this species from Greece (Rhodos and Kos Islands) and Israel. This record makes it probable that the species will be found at other suitable locations in the Mediterranean region as well, most likely in Syria and Turkey.

Taxonomic position. This species was described as the type species of the genus *Lindosiella* (Kritscher, 1959: 454).

Rosin & Shulov (1961) correctly placed the specimens from Israel (Jerusalem) in the genus *Charinus* Simon 1892, but did not attempt their determination to species. Kraus (1961: 491) exa-

mined the specimens from Rhodos and Israel (Jerusalem and Zipari) and concluded that they belong to the same species, but he left them in the genus *Lindosiella*.

While describing two new species of genus *Charinus* from Brazil, Weygoldt (1972) presented a key of all *Charinus* species, in which he included also *Charinus* (= *Lindosiella*) *ioanniticus*.

Quintero (1986) transferred most of the genera from the family Charontidae Simon, 1892 with the type genus *Charon* Karsch, 1879 into the family Charinidae Quintero, 1986 with the type genus *Charinus* Simon, 1892. He does not mention the genus *Lindosiella*. He placed the genera *Cataeus*, *Charinides*, *Charinus*, *Parachron*, *Phrynicosarax*, *Sarax* and *Tricharinus* in the family Charinidae, and left only the genera *Charon* and *Stygophrynus* in the family Charontidae.

ACKNOWLEDGEMENTS. We thank Vladislav Šejna of Prague for providing the Turkish specimens and Jiří Zídek (Socorro, New Mexico, USA) for help with preparation of this account.

REFERENCES

- KRAUS O. 1961: Charontidae aus Israel, ein zoogeographisch bemerkenswertes Vorkommen (Arach., Pedipalpi-Amblypygi). *Senckenbergiana. Biol. (Frankfurt am Main)*, 42: 491-493.
- KRITSCHER E. 1959: Ergebnisse der von Dr. O. Paget und Dr. E. Kritscher auf Rhodos durchgeföhrten zoologischen Exkursionen. *Ann. Naturhist. Mus. Wien*, 63: 453-457.
- QUINTERO D. 1986: Revision de la Clasificacion de Amblypygidos Pulvinados: Creacion de Subordenes, una Nueva Familia y un Nuevo Genero con Tres Nuevas Especies (Arachnida: Amblypygi). *Proc. IX Int. Congr. Arachnol. Panama*, 1983: 203-212.
- ROSIN R. & SHULOV A. 1960: Representatives of the order Amblypygi (Arachnida) found in Israel. *Bull. Res. Counc. Israel.(Zool.)*, 9B: 167-168.
- WEYGOLDT P. 1972: Charontidae (Amblypygi) aus Brasilien. Beschreibung von zwei neuen Charinus-Arten, mit Anmerkungen zur Entwicklung, Morphologie und Tiergeographie und mit einem Bestimmungsschlüssel für die Gattung Charinus. *Zool. Jb. Syst.*, 99: 107-132.

SOUHRN

V práci je uveden první nález bičovce druhu *Charinus ioanniticus* (Kritscher, 1959) z Turecka. Všech 10 exemplářů bylo nalezeno v noci, v asi 30 metrů dlouhé krasové jeskyni, která se nachází v jižním Turecku na okraji vesnice Çevlik poblíž Samandagi. Jeskyně je vstupním otvorem orientována k moři, od kterého je vzdálena cca 400 metrů. Četnými trhlinami ve stěně se do ní dostává deštová voda a trhliny pravděpodobně spojují jeskyni s rozsáhlým systémem dalších nepřístupných prostor.

Charinus ioanniticus byl poprvé nalezen na řeckém ostrově Rhodos. Později byl hlášen ze dvou lokalit v Izraeli (Jeruzaléma a Zipari) a dalšího řeckého ostrova (Kos). S ohledem na zmíněné nálezy je pravděpodobný výskyt druhu na dalších vhodných lokalitách v okolí Středozemního moře (především v Sýrii a Turecku).

Druh byl popsán jako typový druh rodu *Lindosiella* (Kritscher, 1959: 454). Weygoldt (1972) uvádí klíč druhů rodu *Charinus*, kam zařazuje také *Charinus* (= *Lindosiella*) *ioanniticus*. Quintero (1986) přeřazuje většinu rodů z čeledi Charontidae s typovým rodem *Charon* do čeledi Charinidae s typovým rodem *Charinus*. Rod *Lindosiella* neuvádí vůbec. Do čeledi Charinidae řadí rody *Cataeus*, *Charinides*, *Charinus*, *Parachron*, *Phrynicosarax*, *Sarax* a *Tricharinus*. V čeledi Charontidae ponechává pouze rody *Charon* a *Stygophrynus*.